ELSEVIER

Environmental Pollution 130 (2004) I–III

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 130

Number 1 2004

1 Effects of air pollution on the Central and Eastern European mountain forests
A. Bytnerowicz, R. Musselman, R. Szaro

5 Ambient ozone in forests of the Central and Eastern European mountains
A. Bytnerowicz, B. Godzik, K. Grodzińska, W. Frączek, R. Musselman, W. Manning, O. Badea, F. Popescu, P. Fleischer
Ozone concentrations in Central and Eastern European mountain ranges are elevated and phytotoxic to sensitive vegetation.

17 Vegetation of the selected forest stands and land use in the Carpathian Mountains
K. Grodzińska, B. Godzik, W. Frączek, O. Badea, J. Oszlányi, D. Postelnicu, Y. Shparyk
Vegetation and land use maps of forested mountain areas in central Europe are presented.

33 Bioindicator plants for ambient ozone in Central and Eastern Europe
W.J. Manning, B. Godzik
Sixteen species of verified bioindicator plants for ambient ozone are available for use in Central and Eastern Europe.

41 Chemical and morphological characteristics of key tree species of the Carpathian Mountains
B. Maňkovská, B. Godzik, O. Badea, Y. Shparyk, P. Moravčík
Chemical composition of foliage and structure of epicuticular waxes indicated phytotoxic effects of air pollution in many forest sites of the Carpathian Mountains.

55 Heavy metal pollution and forest health in the Ukrainian Carpathians
Y.S. Shparyk, V.I. Parpan
Local industrial emissions of heavy metal pollution and the condition of Ukrainian Carpathians forests are examined.

65 The influence of O_3, NO_2 and SO_2 on growth of *Picea abies* and *Fagus sylvatica* in the Carpathian Mountains
R.M. Muzika, R.P. Guyette, T. Zielonka, A.M. Liebhold
Trends in tree growth rates can be explained by long-term effects of air pollution.

73 Occurrence of spruce bark beetles in forest stands at different levels of air pollution stress
W. Grodzki, M. McManus, M. Knížek, V. Meshkova, V. Mihalciuc, J. Novotny, M. Turčani, Y. Slobodyan
Air pollution (ozone) can be one of predisposing factors that increases the susceptibility of mountain Norway spruce stands to attack by Ips typographus *and associated bark beetle species.*

85 Genetic effects of air pollution on forest tree species of the Carpathian Mountains
R. Longauer, D. Gömöry, L. Paule, I. Blada, F. Popescu, B. Mankovska, G. Müller-Starck, R. Schubert, K. Percy, R.C. Szaro, D.F. Karnosky
Genetic effects of air pollution on main forest trees of the Carpathians are species- and site-specific.

93 Forest health status in the Carpahian Mountains over the period 1997–2001
O. Badea, M. Tanase, J. Georgeta, L. Anisoara, A. Peiov, H. Uhlirova, J. Pajtik, J. Wawrzoniak, Y. Shparyk
About 1/3 of the Carpathian forest trees are damaged by natural stressors and air pollution.

99 Evaluation of air pollution-related risks for Austrian mountain forests
S. Smidt, F. Herman
Despite strong reduction of emissions in Europe, pollutants are still a potential stress factor, especially for sensitive mountain forest ecosystems in Austria.

113 Air pollution and forest health: toward new monitoring concepts
K.E. Percy, M. Ferretti
There is a need for reappraisal of monitoring methods used to assess air pollution effects on forest health.

127 Nature conservation in Central and Eastern Europe with a special emphasis on the Carpathian Mountains
J. Oszlányi, K. Grodzińska, O. Badea, Y. Shparyk
Current status of nature protection in countries of the Carpathian Region is described.

S0269–7491(04)00173–3

Number 2 2004

135 Morphology, chemistry and distribution of neoformed spherulites in agricultural land affected by metallurgical point-source pollution
S. Leguédois, F. van Oort, T. Jongmans, P. Chevallier
Metal spherulites may act as high metal-trapping mineral phases in polluted agricultural soils.

149 Assessment of macro and microelement accumulation capability of two aquatic plants
D. Baldantoni, A. Alfani, P. Di Tommasi, G. Bartoli, A. Virzo De Santo
Element accumulation in roots and shoots of aquatic plants was used as a criteria for selecting useful biomonitors.

157 Effect of anions and cations on cadmium sorption kinetics from aqueous solutions by chitin: experimental studies and modeling
H. Benaissa, B. Benguella
Various ions can influence the ability of chitin to remove cadmium from aqueous solutions.

165 Trichloroacetic acid cycling in Sitka spruce saplings and effects on sapling health following long term exposure
C.A. Dickey, K.V. Heal, R.T. Stidson, R. Koren, P. Schröder, J.N. Cape, M.R. Heal
TCA stored in Sitka spruce needles may affect the health of saplings.

177 Organotin compounds in precipitation, fog and soils of a forested ecosystem in Germany
J.-H. Huang, D. Schwesig, E. Matzner
Forested soils may act as sinks for atmospherically deposited organotin compounds.

187 Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967–2000
A. Kierkegaard, A. Bignert, U. Sellström, M. Olsson, L. Asplund, B. Jansson, C.A. de Wit
PBDEs and MeO-BDEs show different temporal trends in pike from a Swedish lake suggesting different sources.

199 Responses of hybrid poplar clones and red maple seedlings to ambient O_3 under differing light within a mixed hardwood forest
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Under natural forest conditions and ambient O_3 exposures, available light plays a significant role in determining O_3 uptake and resulting physiological changes leading to the development of foliar symptoms.

215 Influence of light fleck and low light on foliar injury and physiological responses of two hybrid poplar clones to ozone
C. Wei, J.M. Skelly, S.P. Pennypacker, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis
Intermittent light flecks enhance physiological injury leading to increased foliar symptoms in clones of ozone-sensitive hybrid poplar.

229 Tracing sewage pollution using linear alkylbenzenes (LABs) in surface sediments at the south end of the Southern California Bight
J.V. Macías-Zamora, N. Ramírez-Alvarez
Linear alkylbenzenes were used to trace the distribution of sewage pollution of surface marine sediments.

239 Sediment trace metal profiles in lakes of Killarney Park, Canada from regional to continental influence
N. Belzile, Y.-W. Chen, J.M. Gunn, S.S. Dixit
Lakes in Killarney Park are in transition from being impacted by regional pollution to being controlled by continental atmospheric deposition.

249 Occurrence of PAHs, PCBs and organochlorine pesticides in the Tonghui River of Beijing, China
Z. Zhang, J. Huang, G. Yu, H. Hong
Organic pollutants were higher in sediments than in surface waters.

263 Macroinvertebrate response to acid mine drainage: community metrics and on-line behavioural toxicity bioassay
A. Gerhardt, L. Janssens de Bisthoven, A.M.V.M. Soares
Toxicity of acid mine drainage was evaluated by macroinvertebrate bioassessment and a new on-line rapid behavioural toxicity test with Atyaephyra desmaresti *(Crustacea).*

275 Effect of cropping and tillage on the dissipation of PAH contamination in soil
C. Saison, C. Perrin-Ganier, M. Schiavon, J.-L. Morel
Dissipation rates of PAHs were slow using natural attentuation.

287 Scale-dependence of land use effects on water quality of streams in agricultural catchments
O. Buck, D.K. Niyogi, C.R. Townsend
Land use influences water quality of streams at various spatial scales.

Number 3 2004

301 Effects of incubation on solubility and mobility of trace metals in two contaminated soils
L.Q. Ma, Y. Dong
Iron is important in controlling metal solubility and mobility in flooded soils.

309 Multi-generation cadmium acclimation and tolerance in *Daphnia magna* Straus
B.T.A. Muyssen, C.R. Janssen
Multi-generation acclimation studies are important for evaluating long-term effects of aquatic pollutants.

317 The role of glomalin, a protein produced by arbuscular mycorrhizal fungi, in sequestering potentially toxic elements
M.C. González-Chávez, R. Carrillo-González, S.F. Wright, K.A. Nichols
Glomalin may be useful in remediation of toxic elements in soils.

325 Natural attenuation of trichloroethene and its degradation products at a lake-shore site
Y.-J. An, D.H. Kampbell, J.W. Weaver, J.T. Wilson, S.-W. Jeong
Field investigation of TCE contamination at a lake-shore site indicates that TCE is anaerobically degrading under ambient conditions.

337 Long-term effects of submergence and wetland vegetation on metals in a 90-year old abandoned Pb–Zn mine tailings pond
D.L. Jacob, M.L. Otte
Metal and acid-volatile sulfide concentrations were affected differently by flooding vegetation.

347 Contamination of butyltin compounds in Malaysian marine environments
A. Sudaryanto, S. Takahashi, H. Iwata, S. Tanabe, A. Ismail
Lack of any regulation of TBT clearly resulted in a heavy contamination of BTs in Malaysia.

359 The influence of season and leaf age on concentrations of radiocaesium (^{137}Cs), stable caesium (^{133}Cs) and potassium in *Agrostis capillaris*
C.A. Salt, J.W. Kay, K.E. Jarvis
Season and leaf age have the same influence on radiocaesium and stable caesium in Agrostis capillaris *but affect K differently.*

371 Haloacetic acids in the aquatic environment. Part I: macrophyte toxicity
M.L. Hanson, K.R. Solomon
Monochloroacetic acid was the most toxic form to aquatic macrophytes.

385 Haloacetic acids in the aquatic environment. Part II: ecological risk assessment
M.L. Hanson, K.R. Solomon
Current environmental concentrations of haloacetic acids do not pose a risk to aquatic macrophytes, but could impact plankton.

403 Comparing aquatic risk assessment methods for the photosynthesis-inhibiting herbicides metribuzin and metamitron
T.C.M. Brock, S.J.H. Crum, J.W. Deneer, F. Heimbach, R.M.M. Roijackers, J.A. Sinkeldam
Laboratory toxicity data for aquatic primary producers can be used to derive ecological thresholds for herbicides.

427 Arbuscular mycorrhizal fungi can decrease the uptake of uranium by subterranean clover grown at high levels of uranium in soil
G. Rufyikiri, L. Huysmans, J. Wannijn, M. Van Hees, C. Leyval, I. Jakobsen
Plant mycorrhization may decrease U concentration in shoots of plants grown at high level of U in soil.

437 Degradation of toxaphene in water during anaerobic and aerobic conditions
M. LacayoR, B. van Bavel, B. Mattiasson
The majority of toxaphene degradation occured under anaerobic conditions.

445 Atmospheric lead deposition to Okefenokee Swamp, Georgia, USA
B.P. Jackson, P.V. Winger, P.J. Lasier
Coal combustion emissions appear to be a major source of Pb in the Okefenokee wetland.

453 Responses of three grass species to creosote during phytoremediation
X.-D. Huang, Y. El-Alawi, D.M. Penrose, B.R. Glick, B.M. Greenberg
Plant growth promoting rhizobacteria enhanced growth and remediation of three grass species.

465 A multi-process phytoremediation system for removal of polycyclic aromatic hydrocarbons from contaminated soils
X.-D. Huang, Y. El-Alawi, D.M. Penrose, B.R. Glick, B.M. Greenberg
Persistent PAH contaminants in soils can be removed more completely and rapidly by using multiple remediation processes.

I Contents of Volume 130